

Chapter 13.14: GLM 3: Factorial designs Factorial ANOVA

Formatted: Font: (Default) Trebuchet MS, 20 pt, English (United Kingdom)

Labcoat Leni's Real Research

Going out on the pierce

Problem

Guéguen, N. (2012). *Alcoholism: Clinical and Experimental Research*, 36(7), 1253–1256.



Tattoos and body piercings have become very popular since I was young. I have often contemplated having Ronald Fisher's face tattooed over my own so that people will think I'm a genius. But I digress. Research has shown that people who have tattoos and piercings are more likely to engage in risky behaviour. Nicolas Guéguen (2012) measured the level of intoxication (mass of alcohol per litre of breath

exhaled, **Alcohol**) in 1965 French youths as they left bars. This measure was an indicator of risky behaviour. Each youth was also classified as having tattoos, piercings, both or neither (**Group**), and their gender was noted (**Gender**). The data are in the file **Gueguen (2012).sav**. Was the level of risk (i.e., alcohol) greater in groups who had tattoos and piercings? Did this effect interact with gender? Draw an error bar chart of the data too.

Solution

To do an error bar chart for means that are independent (i.e., have come from different groups) we need to double-click on the clustered bar chart icon in the Chart Builder (see the book chapter). All we need to do is to drag our variables into the appropriate drop zones. Select **Alcohol** from the variable list and drag it into ; select **Group** from the variable list and drag it into ; finally, select the **Gender** variable and drag it into . This will mean that lines representing males and females will be displayed in different colours. Select error bars in the properties dialog box and click on to apply them to the Chart Builder. Click on to produce the graph.




Figure 1

Figure 1 is the resulting error bar graph of these data. Looking at the graph, we can see that in each group the men had consumed more alcohol than the women (the blue bars are taller than the green bars for all groups); this suggests that there may be a significant main effect of **Gender**. There is a steady increase in the volume of alcohol consumed as we move along the **Group** variable – the *no tattoos, no piercing* group consumed the least amount of alcohol and the *tattoos and piercings* group consumed the largest amount of alcohol – suggesting that there may be a significant main effect of **Group**. This trend appears to be the same for both men and women, suggesting that the interaction effect of **Gender** and **Group** is unlikely to be significant.

We need to conduct a 4 (experimental group) \times 2 (gender) two-way independent ANOVA on the mass of alcohol per litre of exhaled breath. To access the main dialog box for a general factorial ANOVA select **Analyze** **General Linear Model** **Univariate...**. First, select the dependent variable **Alcohol** from the variables list on the left-hand side of the dialog box and drag it to the space labelled *Dependent Variable*. In the space labelled *Fixed Factor(s)* we need to place any independent variables relevant to the analysis. Select **Group** and **Gender** in the variables list (these variables can be selected simultaneously by holding down *Ctrl* while clicking on the variables) and drag them to the *Fixed Factor(s)* box.

DISCOVERING STATISTICS USING SPSS

Descriptives

Gender		Statistic	Std. Error	
Mass of alcohol per liter of exhaled breath	Male	Mean	.1909	
		95% Confidence Interval for Mean	Lower Bound	.1822
			Upper Bound	.1995
		5% Trimmed Mean	.1827	
		Median	.1700	
		Variance	.021	
		Std. Deviation	.14544	
		Minimum	.00	
		Maximum	.81	
		Range	.81	
		Interquartile Range	.14	
		Skewness	.790	.074
		Kurtosis	.202	.149
		Female	Mean	.1495
	95% Confidence Interval for Mean		Lower Bound	.1422
			Upper Bound	.1569
	5% Trimmed Mean		.1408	
Median	.1500			
Variance	.012			
Std. Deviation	.11101			
Minimum	.00			
Maximum	.66			
Range	.66			
Interquartile Range	.14			
Skewness	1.202	.082		
Kurtosis	2.854	.164		

Output 1

Tests of Between-Subjects Effects

Dependent Variable: Mass of alcohol per liter of exhaled breath

Source	Type III Sum of Squares	df	Mean Square	F	Sig.	Partial Eta Squared
Model	60.869 ^a	8	7.609	465.174	.000	.655
Gender	.269	1	.269	16.443	.000	.008
Group	1.319	3	.440	26.883	.000	.040
Gender * Group	.080	3	.027	1.624	.182	.002
Error	32.010	1957	.016			
Total	92.879	1965				

a. R Squared = .655 (Adjusted R Squared = .654)

Output 2

Multiple Comparisons

Dependent Variable: Mass of alcohol per liter of exhaled breath
LSD

(I) Group	(J) Group	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
No Tattoo, No Piercing	Tattoos Only	.0039	.01019	.704	-.0161	.0238
	Piercings Only	-.0522*	.00898	.000	-.0698	-.0345
	Tattoos and Piercings	-.0805*	.01255	.000	-.1051	-.0558
Tattoos Only	No Tattoo, No Piercing	-.0039	.01019	.704	-.0238	.0161
	Piercings Only	-.0560*	.01272	.000	-.0810	-.0311
	Tattoos and Piercings	-.0843*	.01544	.000	-.1146	-.0540
Piercings Only	No Tattoo, No Piercing	.0522*	.00898	.000	.0345	.0698
	Tattoos Only	.0560*	.01272	.000	.0311	.0810
	Tattoos and Piercings	-.0283	.01467	.054	-.0571	.0005
Tattoos and Piercings	No Tattoo, No Piercing	.0805*	.01255	.000	.0558	.1051
	Tattoos Only	.0843*	.01544	.000	.0540	.1146
	Piercings Only	.0283	.01467	.054	-.0005	.0571

Based on observed means.

The error term is Mean Square(Error) = .016.

*. The mean difference is significant at the .05 level.

Output 3

3. Group

Dependent Variable: Mass of alcohol per liter of exhaled breath

Group	Mean	Std. Error	95% Confidence Interval	
			Lower Bound	Upper Bound
No Tattoo, No Piercing	.154	.003	.147	.161
Tattoos Only	.168	.010	.148	.189
Piercings Only	.216	.008	.200	.233
Tattoos and Piercings	.248	.014	.220	.276

Output 4

[Output Output-2](#) is the output of the main ANOVA. We can see that there was a significant main effect of gender, $F(1, 1957) = 16.44, p < .001$, with a partial eta squared of .01. The means in [Output Output-1](#) (and [Figure Figure-1](#)), reveal that men ($M = 0.19, SD = 0.15$) consumed a significantly higher mass of alcohol than women ($M = 0.15, SD = 0.11$). There was also a significant main effect of group, $F(3, 1957) = 26.88, p < .001$, with a partial eta squared of .04. *Post hoc* tests ([Output Output-3](#)) revealed that participants who only had piercings ($M = 0.22$) consumed a significantly greater mass of alcohol than those who only had tattoos ($M = 0.17$) (least significant difference (LSD) test, $p < .001$) and those who had no tattoos and no piercings ($M = 0.15$) (LSD test, $p < .001$). Participants who had both tattoos and piercings ($M = 0.25$) consumed a significantly greater mass of alcohol than those who only had tattoos ($M = 0.17$) (LSD test, $p < .001$), and those who had no tattoos and no piercings ($M = 0.15$) (LSD test, $p < .001$). However, they did not consume a significantly greater mass than those who only had piercings ($M = 0.22$) (LSD test, $p = .05$).

DISCOVERING STATISTICS USING SPSS

In summary, we can conclude that individuals who have both piercings and tattoos, and those who only have piercings, consumed significantly more alcohol than those who had no tattoos and no piercings and those who only had tattoos. This effect was found in both men and women.

~~Don't forget your toothbrush?~~